

## EVALUATION OF *Plasmodium cynomolgi* SPOROZOITES INDUCED INFECTIONS OF CAPTIVE BORN *Macaca Fascicularis*

Principal Investigators : George S. Ward, LTC, VC  
Pranee Hansukjariya, BSc.  
Suwatana Wongsepradit  
Richard G. Andre, MAJ(P) MSC  
David E. Davidson, Jr., COL, VC

**OBJECTIVE :** To determine if the cynomolgus monkey can be used to supplement rhesus monkeys in the antimalarial radical curative test.

**BACKGROUND :** India has ceased exportation of rhesus monkeys which are used in the *Plasmodium cynomolgi* antimalarial compound testing model. The cynomolgus (*Macaca fascicularis*) is raised at AFRIMS and this species is more productive in a laboratory breeding colony than are rhesus monkeys. A systematic evaluation of captive born has not been completed to determine if this species could be used to supplement scarce rhesus monkeys.

**RESULTS :** Twenty-five AFRIMS produced cynomolgus have been inoculated with *P. cynomolgi* sporozoites. The intact cynomolgus does not develop an infection comparable to the rhesus. The splenectomized captive-born cynomolgus monkey appears to be capable of supplementing rhesus in antimalarial compound testing. Relapse occurs (as in rhesus monkeys) after clearance of blood forms with chloroquine given alone or alone with a noncurative dose of primaquine or test compound.

**FUTURE OBJECTIVES :** To date only 7 splenectomized cynomolgus have had evaluation of relapse potential following clearance of blood forms. This number needs to be expanded to allow comparison with intact rhesus. The dose of sporozoites injected into splenectomized cynomolgus needs to be standardized so future test animals will be inoculated with the same dose as the rhesus in the antimalarial agent testing project.

### REFERENCES :

1. Kinnamon, K.E., and Rothe, W.E., 1973. Biological screening in the U.S. Army antimalarial drug development program *Am. J. Trop. Med. Hyg.* 24: 174-178.
2. Schmidt, L.H.I., Fradkin, R., Vaughan, D., and Rasco, J., 1977. Radical cure of infections with *Plasmodium cynomolgi*: A function of total 8-aminoquinoline dose. *Am. J. Trop. Med. Hyg.*, 26: 1116-1128.
3. Davidson, D.E., Jr., Agerr, A.L., Brown, J.L., Chapple, F.E., Whitmire, R.E., and Rossan, R.N. 1981. New tissue schizonticidal antimalarial drugs. *Bull WHO* 59(3): 463-79.

4. Krootoski, W.A., Collins, W.E., Broderson, J.R., Warren, M., and Krotoski, D.M., 1981. The 48-hour exoerythrocytic stage of *Plasmodium cynomolgi bastianellii*. Am. J. Trop. Med. Hyg. 30: 31-37.
5. Schmidt, L.H., Gramer, D.V., Rossan, R.N., and Harrison, J., 1977. The characteristics of *Plasmocium cynomolgi* infections in various old world primates. Am. J. Trop. Med. Hyg. 26: 356-372.
6. Coatney, G.R., Collins, W.E., Warren, M. and Contacos, P.G. 1971. *The Primate Malariae* U.S. Gov. Printing Office, Washington, D.C.
7. Schmidt, L.H., Rossan, R.N., and Fisher, K.P., 1963. The activity of a repository form of 4, 6-diamino-1-(p-xhloeophwny)-1-2-sihyseo-2, 2-simwrhyl-5-triInw against infections with *Plasmodium cynomolgi*. Am. J. Trop. Med. Hyg., 12: 494-503.
8. Davidson, D.E., Johnson, D.O., Tanticharoenyos, P., Hickman, R.L. and Kinnon, K.E., 1976. Evaluating new antimalarial drugs against trophozoite induced *Plasmocium cynomolgi* malaria in rhesus monkeys. Am. J. Trop. Med. Hyg. 25: 26-33.
9. Schmidt, L.H., Pradkin, R., Genther, C.S., Rossana, R.N. and Squires, W., 1982. The characterisitics of untreated sporozoite-induced and trophozoite-induced infections. Am. J. Trop. Hyg...., 31: 612-645.